arXiv:1702.03084v1 [math.DS] 10 Feb 2017

MEASURE RIGIDITY FOR SOLVABLE GROUP ACTIONS IN THE SPACE OF LATTICES

MANFRED EINSIEDLER AND RONGGANG SHI

Abstract. We study invariant probability measures on the homogeneous space $\mathrm{SL}_n(\mathbb{R})/\mathrm{SL}_n(\mathbb{Z})$ for the action of subgroups of $\mathrm{SL}_n(\mathbb{R})$ of the form SF where F is generated by one parameter unipotent groups and S is a one parameter $\mathbb{R}$-diagonalizable group normalizing F. Under the assumption that S contains an element with only one eigenvalue less than one (counted with multiplicity) and others bigger than one we prove that all the SF invariant and ergodic probability measures on $\mathrm{SL}_n(\mathbb{R})/\mathrm{SL}_n(\mathbb{Z})$ are homogeneous.

1. introduction

Let $n \geqslant 2$ be a positive integer. Let $G = \mathrm{SL}_n(\mathbb{R}), \Gamma = \mathrm{SL}_n(\mathbb{Z})$ and $X = G/\Gamma$. Let S be a one parameter $\mathbb{R}$-diagonalizable subgroup of G, i.e. S is conjugate in G to a subgroup with diagonal matrices. Let F be a subgroup of G normalized by S and generated by one parameter unipotent subgroups. This paper is about the classification of SF invariant and ergodic probability measures on X.

This type of questions has already been studied earlier by Margulis-Tomanov [4] and Mozes [5] using Ratner's measure rigidity theorem [7] for F actions. In particular, it is proved in [4] and [5] that if SF is an epimorphic subgroup of $H \leqslant G$ and H is generated by one parameter unipotent subgroups, then any SF invariant and ergodic probability measure is H invariant and hence homogeneous by Ratner's theorem. Recently, in a joint paper with Barak Weiss the second named author [8] proved measure rigidity for certain non-epimorphic sovable subgroup actions on X for $n = 3$. More precisely, it is proved in [8, Theorem 5] that if SF is nonabelian and S has no nonzero fixed vectors in $\mathbb{R}^3$, then SF acts uniquely ergodically on X. It is noticed already in [8] that even for $n = 4$, there are homogeneous and even non-homogeneous SF

2000 *Mathematics Subject Classification.* Primary 37C85; Secondary 28A33, 22E40.

Key words and phrases. homogeneous dynamics, measure rigidity.

R. Shi is supported by NSFC (11201388) and Fundamental Research Funds for the Central Universities (20720160006).

invariant and ergodic probability measures under similar assumptions. Therefore, to get rigidity results one needs to put further restrictions on the group S.

A one parameter $\mathbb{R}$-diagonalizable subgroup of G is said to be simple if it is conjugate to

$$\{\mathrm{diag}(e^{r_1 t}, \dots, e^{r_{n-1} t}, e^{-t}) : t \in \mathbb{R}\}$$

for a probability vector $(r_1, \dots, r_{n-1}) \in \mathbb{R}^{n-1}$ with positive entries. The following is the main result of this paper.

Theorem 1.1. *Let F be a subgroup of $G = \mathrm{SL}_n(\mathbb{R})$ $(n \geqslant 2)$ generated by one parameter unipotent subgroups. Let $S = \{a_t : t \in \mathbb{R}\}$ be a simple one parameter $\mathbb{R}$-diagonalizable subgroup of G normalizing F. Then any SF invariant and ergodic probability measure on $X = G/\Gamma$ $(\Gamma = \mathrm{SL}_n(\mathbb{Z}))$ is homogeneous.*

The starting point of the proof of Theorem 1.1 is the same as that of [4] and [5] where μ is written as a convex comination of F invariant and ergodic probability measures. Note that any F invariant and ergodic probability measure ν on X is homogeneous by Ratner's theorem, i.e. there is a closed subgroup H of G and $x_g \in X$ (here x_g is the coset $g\Gamma$ for $g \in G$) such that ν is supported on Hx_g and is H invariant. We show that $a_t\nu$ converges to zero in the space of finite measures on X as t tends to infinity or minus infinity unless $g^{-1}Hg$ is a semisimple algebraic group virtually defined over $\mathbb{Q}$ and its action on $\mathbb{R}^n$ has only one $\mathbb{Q}$-isotropic type. The main new ingredient here is the use of Tits [9] on the structure of irreducible representations to show that the centralizer of H in G (denoted by $Z_G(H)$) is an abelian group. The proof is then completed by a standard argument.

Acknowledgements. The authors would like to thank MSRI for its hospitality during Spring 2015.

2. Irreducible representations over $\mathbb{Q}$

This section is about the structure of $\mathbb{Q}$-rational representations of semisimple algebraic groups defined over $\mathbb{Q}$. We adopt the convention of [1] to identify algebraic varieties or groups with their rational points over complex numbers and denote them by boldfaced capital letters.

Let $\mathbf{V}$ be an n-dimensional vector space over $\mathbb{C}$ endowed with a $\mathbb{Q}$-structure. Let $\mathbf{G} = \mathrm{SL}(\mathbf{V})$ be the special linear group with the natural $\mathbb{Q}$-structure such that linear action of $\mathbf{G}$ on $\mathbf{V}$ is a $\mathbb{Q}$-rational representation. Let $\mathbf{H}$ be nontrivial connected semisimple subgroup of $\mathbf{G}$ defined over $\mathbb{Q}$. Recall that the linear action of $\mathbf{H}$ on $\mathbf{V}$ is completely

determined by the induced Lie algebra representations which is completely reducible over $\mathbb{Q}$ (see [13, Theorem 7.8.11]). So $\mathbf{V}$ is a direct sum of $\mathbf{H}$ invariant $\mathbb{Q}$-irreducible subspaces. We say the linear action of $\mathbf{H}$ on $\mathbf{V}$ has only one $\mathbb{Q}$-isotropic type if all these irreducible subspaces are isomorphic to each other over $\mathbb{Q}$.

For a field $\mathbb{F}$ containing $\mathbb{Q}$ and a variety $\mathbf{X}$ defined over $\mathbb{Q}$, we let $\mathbf{X}_{\mathbb{F}}$ be the set of $\mathbb{F}$-rational points of $\mathbf{X}$. Let $V = \mathbf{V}_{\mathbb{R}}$, $G = \mathbf{G}_{\mathbb{R}}$ and $H = \mathbf{H}_{\mathbb{R}}$. The main result of this section is the following proposition.

Proposition 2.1. *Let S be a simple one parameter $\mathbb{R}$-diagonalizable subgroup of G. Suppose S is contained in $HZ_G(H)$ and the linear action of $\mathbf{H}$ on $\mathbf{V}$ has only one $\mathbb{Q}$-isotropic type. Then $Z_G(H)$ is an abelian group.*

Proof. Since the linear action of $\mathbf{H}$ on $\mathbf{V}$ has only one $\mathbb{Q}$-isotropic type, there exists a positive integer m such that $\mathbf{V}$ is isomorphic to the direct sum of m copies of a certain $\mathbb{Q}$-irreducible representation of $\mathbf{H}$. According to [9, Theorem 7.2], there exists a central simple division algebra $\mathbb{D}$ over a number field $\mathbb{F}$, a complex vector space $\mathbf{W}$ defined over $\mathbb{F}$ with $\mathbf{W}_{\mathbb{F}}$ having a right $\mathbb{D}$-module structure and an absolutely irreducible $(\mathbb{D}, \mathbb{F})$-representation $\rho : \mathbf{H} \to \mathrm{GL}(\mathbf{W})$ such that

$$\mathbf{V} \cong \mathrm{Res}_{\mathbb{F}/\mathbb{Q}}\big(\mathrm{Res}_{\mathbb{D}/\mathbb{F}}\mathbf{W}^{\oplus m}\big) \tag{2.1}$$

in the category of $\mathbb{Q}$-rational representations of $\mathbf{H}$. According to Tits' notation, the morphism ρ is defined over $\mathbb{F}$ and $\rho(\mathbf{H}_{\mathbb{F}})$ acts $\mathbb{D}$-linear on $\mathbf{W}_{\mathbb{F}}$. The notion of absolute irreducibility will be explained later.

Let $d^2 = [\mathbb{D} : \mathbb{F}]$ and $k = [\mathbf{W}_{\mathbb{F}} : \mathbb{D}]$. We will show that $m = d = 1$, which will imply the proposition immediately. We would like to understand the group H and its centralizers by choosing a good basis of V using (2.1). To make the presentation simpler we assume the left and right hand sides of (2.1) are equal. We will give explicit descriptions of $\mathbb{D}, \mathbf{W}$ and of restriction of scalars using several isomorphisms.

By choosing an isomorphism $\mathbb{D}\otimes_{\mathbb{F}}\mathbb{C} \cong \mathrm{M}_d(\mathbb{C})$ and identifying $\mathbb{D}$ with $\mathbb{D} \otimes 1$ in the natural way we assume that $\mathbb{D}$ is a subring of $\mathrm{M}_d(\mathbb{C})$. We assume $\mathbf{W}_{\mathbb{F}} = \mathbb{D}^k$ and $\mathbf{W} = \mathrm{M}_d(\mathbb{C})^k$ where $\mathrm{M}_d(\mathbb{C})^k$ is naturally identified with the set of kd-by-d matrices by viewing its elements blockwise. Therefore, the group of right $\mathrm{M}_d(\mathbb{C})$-module automorphisms of $\mathbf{W}$ can be naturally identified with $\mathrm{GL}_{kd}(\mathbb{C})$ which acts on $\mathbf{W}$ from the left. Let $\widehat{\rho}$ be the composite of ρ and this identification, that is, $\widehat{\rho}(\mathbf{H})$ is a subgroup of $\mathrm{GL}_{kd}(\mathbb{C})$. We consider $\widehat{\rho}$ as a morphism of algebraic groups defined over $\mathbb{F}$ with

$$\widehat{\rho}(\mathbf{H}_{\mathbb{F}}) \leqslant \mathrm{GL}_k(\mathbb{D}).$$

We note that the absolute irreducibility of ρ refers to the fact that $\widehat{\rho}(\mathbf{H})$ acts irreducibly on $\mathbb{C}^{kd}$.

Let $\sigma_1, \dots, \sigma_r$ be the real embeddings of $\mathbb{F}$ and $\sigma_{r+1}, \sigma'_{r+1}, \dots, \sigma_{r+s}, \sigma'_{r+s}$ be the pairs of complex embeddings of $\mathbb{F}$. Each σ_i defines a functor from the category of varieties defined over $\mathbb{F}$ to the category of varieties defined over $\mathbb{F}^{\sigma_i} \stackrel{\text{def}}{=} \{\sigma_i(x) : x \in \mathbb{F}\}$ (see [1, §0.14.1] for definitions). For a variety $\mathbf{X}$ defined over $\mathbb{F}$ we let $\mathbf{X}^{\sigma_i}$ be the image of this functor. Also let $\mathbb{D}^{\sigma_i} = \mathbb{D} \otimes_{\mathbb{F}} \mathbb{F}^{\sigma_i}$ where the left $\mathbb{F}$-module structure of $\mathbb{F}^{\sigma_i}$ comes from the left multiplication of $\sigma_i(\cdot)$.

According to (2.1) and [12, Theorem 1.3.2] we have

$$V = \prod_{i=1}^{r+s} V_i \quad \text{where} \quad V_i = \begin{cases} \left(\mathbf{W}^{\sigma_i}_{\mathbb{R}}\right)^{\oplus m} & 1 \leqslant i \leqslant r \\ \left(\mathbf{W}^{\sigma_i}\right)^{\oplus m} & r+1 \leqslant i \leqslant r+s \end{cases}$$

and

$$H = \prod_{i=1}^{r+s} H_i \quad \text{where} \quad H_i = \begin{cases} \rho^{\oplus m}(\mathbf{H})^{\sigma_i}_{\mathbb{R}} & 1 \leqslant i \leqslant r \\ \rho^{\oplus m}(\mathbf{H})^{\sigma_i} & r+1 \leqslant i \leqslant r+s. \end{cases} \tag{2.2}$$

Moreover,

$$Z_G(H) = G \cap \prod_{i=1}^{r+s} Z_i,$$

where Z_i is the centralizer of H_i in $\mathrm{GL}_{\mathbb{R}}(V_i)$. It suffices to prove that all the Z_i are abelian groups.

For $r+1 \leqslant i \leqslant r+s$, the complex part of the above decomposition is easy to understand. Since $\mathbb{D}^{\sigma_i} \otimes_{\mathbb{F}^{\sigma_i}} \mathbb{C} = \mathrm{M}_d(\mathbb{C})$, we have a decomposition

$$V_i = V'_i \otimes_{\mathbb{C}} V''_i$$

where $V'_i = \mathbb{C}^{kd}$ and $V''_i = \mathbb{C}^{md}$ such that the group

$$H_i = \widehat{\rho}(\mathbf{H})^{\sigma_i} \otimes \mathrm{I}_{V''_i} \tag{2.3}$$

where $\mathrm{I}_{V''_i}$ denotes the identity map of V''_i.

For the real part we first note that the central simple algebra $\mathbb{D}^{\sigma_i} \otimes_{\mathbb{F}^{\sigma_i}} \mathbb{R}$ is equal to $\mathrm{M}_d(\mathbb{R})$ or $\mathrm{M}_{d/2}(\mathbb{H})$ where in the latter case $\mathbb{H}$ is the Hamilton quaternions and d has to be an odd number. By choosing an embedding $\mathbb{H} \to \mathrm{M}_2(\mathbb{C})$ we assume that $\mathrm{M}_{d/2}(\mathbb{H})$ is a subring of $\mathrm{M}_d(\mathbb{C})$. Suppose for $1 \leqslant i \leqslant l$ we are in the case of $\mathrm{M}_d(\mathbb{R})$ and for $l+1 \leqslant i \leqslant r$ we are in the case of $\mathrm{M}_{d/2}(\mathbb{H})$. Then

$$V_i = V'_i \otimes_{\mathbb{R}} V''_i \quad \text{and} \quad H_i = \widehat{\rho}(\mathbf{H})^{\sigma_i}_{\mathbb{R}} \otimes \mathrm{I}_{V''_i} \tag{2.4}$$

where

$$V_i' = \mathbb{R}^{kd} \text{ and } V_i'' = \mathbb{R}^{md} \quad \text{for } 1 \leqslant i \leqslant l;$$
$$V_i' = \mathbb{H}^{kd/2} \text{ and } V_i'' = \mathbb{R}^{md/2} \quad \text{for } l+1 \leqslant i \leqslant r.$$

For $1 \leqslant i \leqslant r$ (resp. $r+1 \leqslant i \leqslant r+s$) let R_i be the ring of $\mathbb{R}$-linear spans of $\widehat{\rho}(\mathbf{H})_{\mathbb{R}}^{\sigma_i}$ (resp. $\widehat{\rho}(\mathbf{H})^{\sigma_i}$) in the ring of $\mathbb{R}$-linear transformations of V_i'. Since $\widehat{\rho}(\mathbf{H})$ acts irreducibly on $\mathbb{C}^{kd}$, the space V_i' is an irreducible R_i-module. For every i there is a division algebra D_i such that the opposite division algebra D_i° is isomorphic to set of R_i-module morphisms of V_i'. By Wedderburn-Artin theorem (see e.g. [3, §1]), the ring R_i is isomorphic to a matrix ring over D_i and V_i is isomorphic to a free D_i module of finite rank. Since $\widehat{\rho}(\mathbf{H})$ acts irreducibly on $\mathbb{C}^{kd}$, we have

$$D_i = \begin{cases} \mathbb{R} & 1 \leqslant i \leqslant l \\ \mathbb{H} & l+1 \leqslant i \leqslant r \\ \mathbb{C} & r+1 \leqslant i \leqslant r+s. \end{cases} \tag{2.5}$$

We can understand Z_i using (2.5), (2.3) and (2.4). For $r+1 \leqslant i \leqslant r+s$,

$$Z_i = (\mathbb{C}\mathrm{I}_{V_i'} \otimes_{\mathbb{C}} \mathrm{Hom}_{\mathbb{C}}(V_i'', V_i''))^* = \mathrm{I}_{V_i'} \otimes \mathrm{GL}_{\mathbb{C}}(V_i'')$$

where $(\cdot)^*$ denotes the invertible elements of a ring. Moreover, if $c_i \in Z_i$ is diagonalizable over $\mathbb{R}$, then we have a decomposition $c_i = \mathrm{I}_{V_i'} \otimes b_i$ where b_i is an $\mathbb{R}$-diagonal element of $\mathrm{GL}_{\mathbb{C}}(V_i'')$ (here we view $\mathrm{GL}_{\mathbb{C}}(V_i'')$ as the real points of an algebraic group coming from the restriction of scalars). For $1 \leqslant i \leqslant l$

$$Z_i = \left(\mathbb{R}\mathrm{I}_{V_i'} \otimes_{\mathbb{R}} \mathrm{Hom}_{\mathbb{R}}(V_i'', V_i'')\right)^* = \mathrm{I}_{V_i'} \otimes \mathrm{GL}_{\mathbb{R}}(V_i''),$$

and every $\mathbb{R}$-diagonalizable $c_i \in Z_i$ can be decomposed as

$$\mathrm{I}_{V_i'} \otimes b_i$$

where $b_i \in \mathrm{GL}_{\mathbb{R}}(V_i'')$ is $\mathbb{R}$-diagonalizable. For $l+1 \leqslant i \leqslant r$

$$Z_i = \left(\mathbb{H}\mathrm{I}_{V_i'} \otimes_{\mathbb{R}} \mathrm{Hom}_{\mathbb{R}}(V_i'')\right)^* \cong \mathrm{GL}_{md/2}(\mathbb{H}\mathrm{I}_{V_i'}),$$

where $\mathbb{H}\mathrm{I}_{V_i'}$ acts on V_i' via right multiplications of matrices. Suppose $c_i \in Z_i$ is $\mathbb{R}$-diagonalizable, then there exists $g_i \in Z_i$ such that

$$g_i c_i g_i^{-1} = \mathrm{I}_{V_i'} \otimes b_i$$

where $b_i \in \mathrm{GL}_{\mathbb{R}}(V_i'')$ is $\mathbb{R}$-diagonalizable.

Now we turn to the proof of the conclusion. Since S is contained in $HZ_G(H)$, the group S is the direct product S_1S_2 of one parameter $\mathbb{R}$-diagonalizable subgroups $S_1 \leqslant H$ and $S_2 \leqslant Z_G(H)$. Since S is simple, there exists $a \in S$ such that the linear action of a on $\mathbb{R}^n$ has only one eigenvalue $\lambda < 1$ counted with multiplicity and all the other

eigenvalues are bigger than 1. We write $a = hb$ where $h \in S_1$ and $b \in S_2$. According to the discussions previously, by possibly replacing S by gSg^{-1} for some $g \in \prod_{i=1}^{r+s} Z_i$ we have

$$h = \prod_{i=1}^{r+s} h_i \otimes \mathrm{I}_{V_i''} \quad \text{and} \quad b = \prod_{i=1}^{r+s} \mathrm{I}_{V_i'} \otimes b_i \tag{2.6}$$

where all the h_i and b_i are $\mathbb{R}$-diagonalizable elements of $\mathrm{GL}_{\mathbb{R}}(V_i')$ and $\mathrm{GL}_{\mathbb{R}}(V_i'')$ respectively.

There exists i_0 with $1 \leqslant i_0 \leqslant r+s$ such that λ is an eigenvalue of a restricted to V_{i_0}. Since each eigenvalue of a restricted to V_i has multiplicity at least 4 (resp. 2) for $l < i \leqslant r$ (resp. $r+1 \leqslant i \leqslant r+s$), we have $i_0 \leqslant l$. We assume without loss of generality that $i_0 = 1$. Since $\det h_i = 1$ for all i, λ has multiplicity 1, all the other eigenvalues of a are bigger than 1 and (2.6), we have $\det b_i > 1$ for $i > 1$. Since $\det a = 1$ we have $0 < \det b_1 \leqslant 1$ (with equality in case $\mathbb{F} = \mathbb{Q}$). This and the simplicity of λ implies that $\min\{kd, md\} = 1$. Therefore $d = 1$. Since $\mathbf{H}$ is a semisimple algebraic group, we have $k > 1$ and hence $m = 1$. Using previously discussed structures of $Z_G(H)$ and $m = d = 1$, we have $l = r$ and $Z_G(H)$ is an abelian group.

□

3. MEASURE RIGIDITY

The aim of this section is to prove Theorem 1.1. Let G, Γ, X, S, F be as in Theorem 1.1. For every $g \in G$ we use x_g to denote the coset $g\Gamma \in X$. Also, we use x_Γ to denote the coset $\Gamma \in X$. We endow G and $\mathbb{R}^n$ with $\mathbb{Q}$-structures such that their $\mathbb{Q}$-rational points are $\mathrm{SL}_n(\mathbb{Q})$ and $\mathbb{Q}^n$ respectively. A subgroup H of G is said to be virtually defined over $\mathbb{Q}$ if it has finite index in the group of real points of a $\mathbb{Q}$-rational algebraic group. The following is a version of Ratner's theorem [7], see also the work [2, Prop. 1.1] of Borel-Prasad and [10, Theorem 2] of Tomanov.

Theorem 3.1. *Let ν be an F invariant and ergodic probability measure on X. Then there exists a connected subgroup H of G virtually defined over $\mathbb{Q}$ with semisimple Levi factor and a point $x_g \in X$ such that the measure ν is invariant under the action of $gHg^{-1} \geqslant F$ and ν is concentrated on the closed subset gHx_Γ.*

Lemma 3.2. *Suppose there is a proper $\mathbb{Q}$-rational subspace V of $\mathbb{R}^n$ invariant under SH where H is a subgroup of G normalized by S. Then either $\{a_t h x_\Gamma : t \geqslant 0\}$ is divergent for every $h \in H$ or $\{a_t h x_\Gamma : t \leqslant 0\}$ is divergent for every $h \in H$.*

Proof. By assumption, there exists an integer k with $1 \leqslant k \leqslant n-1$ and a basis $v_1, \dots, v_k$ of V consisting of rational vectors. Since the space V is SH invariant, the vector $w = v_1 \wedge \cdots \wedge v_k \in \wedge^k \mathbb{Q}^n$ is an eigenvector of S. By assumption the one parameter group S is simple, so there exists a nonzero real number λ such that $a_t w = e^{\lambda t} w$. Therefore $a_t w \to 0$ as $t \to \infty$ or $t \to -\infty$. We assume without loss of generality that $a_t w \to 0$ as $t \to \infty$. Note that for any $h \in H$ one has $hw = sw$ for some nonzero real number s, which implies that $a_t h w \to 0$ as $t \to \infty$. It follows from the Mahler's compactness criterion (see e.g. [13, Propositon 3.1]) that the trajectory $\{a_t h x_\Gamma : t \geqslant 0\}$ is divergent for every $h \in H$. □

Proof of Theorem 1.1. Let μ be an SF invariant and ergodic probability measure on X. It follows from Poincare recurrence that for almost every $x \in X$

(1) Neither of the trajectories $\{a_t x : t \geqslant 0\}$ or $\{a_t x : t \leqslant 0\}$ is divergent.

Let

$$\mu = \int_X \mu_x \, \mathrm{d}\mu(x)$$

be the ergodic decomposition of μ into F invariant and ergodic components. According to [4] and [5] we can find a connected subgroup H of G and a measurable subset X' of X with full μ measure such that for every $x \in X'$ the following properties hold:

(2) μ_x is H invariant and supported on the closed subset Hx.
(3) $S \leqslant N_G^1(H)^\circ$ where $N_G^1(H)$ consists of $g \in N_G(H)$ with the property that the conjugation by g preserves the Haar measure of H. Moreover, $\mu(N_G^1(H)^\circ x) = 1$.

By possibly removing a measure zero subset from X', we assume that (1) holds for all $x \in X'$.

According to (2) and Theorem 3.1 there is a connected subgroup H' of G virtually defined over $\mathbb{Q}$ with semisimple Levi factor and $g \in G$ with $x_g \in X'$ such that $H = gH'g^{-1}$. By replacing S by $g^{-1}Sg$, F by $g^{-1}Fg$ and μ by $g^{-1}\mu$ we can without loss of generality assume that $H = H'$ and $x_\Gamma \in X'$.

Let $\mathscr{R}_u(H)$ be the maximal unipotent normal subgroup of H. Then H has a connected semisimple subgroup H_0 virtually defined over $\mathbb{Q}$ such that $H = H_0 \ltimes \mathscr{R}_u(H)$. Note that $\Gamma \cap H_0$ and $\Gamma \cap \mathscr{R}_u(H)$ are lattices in H_0 and $\mathscr{R}_u(H)$ respectively. As a connected unipotent group, the space of $\mathscr{R}_u(H)$ invariant vectors

$$V \stackrel{\text{def}}{=} \{v \in \mathbb{R}^n : hv = v \text{ for all } h \in \mathscr{R}_u(H)\}$$

is nonzero and defined over $\mathbb{Q}$. Since S normalizes H, the space V is also S-invariant. If $\mathscr{R}_u(H)$ is nontrivial, then V is a proper subspace of $\mathbb{R}^n$. Hence Lemma 3.2 implies that either $\{a_t x_\Gamma : t \geqslant 0\}$ or $\{a_t x_\Gamma : t \leqslant 0\}$ is divergent. This contradicts (1). Therefore $\mathscr{R}_u(H)$ is trivial and $H = H_0$ is semisimple.

Let $Z = Z_G(H)$. Note that $N_G(H)$ and HZ have the same connected component, see [6]. So by (3) we have $S \leqslant HZ$. The natural representation of H on $\mathbb{R}^n$ is defined over $\mathbb{Q}$ and is completely reducible over $\mathbb{Q}$. Since S is contained in HZ, every maximal $\mathbb{Q}$-rational H-invariant subspace which has only one $\mathbb{Q}$-isotropic type is S-invariant. By Lemma 3.2 and (1) the representation of H on $\mathbb{R}^n$ has only one $\mathbb{Q}$-isotropic type. Therefore Proposition 2.1 implies that Z is an abelian group. In view of (3) we get $\mu(ZHx) = 1$. By pulling μ back to ZH/Γ_1 where $\Gamma_1 = ZH \cap \Gamma$ we get an SF-invariant and ergodic probability measure μ_1 on ZH/Γ_1. Note that SF is generated by Ad-unipotent one parameter subgroups of ZH. It follows from Ratner's theorem that μ_1 is a homogeneous measure. Therefore μ is homogeneous.

□

References

[1] A. Borel, Linear algebraic groups, 2nd ed., GTM 126, Springer-Verlag, 1991.

[2] A. Borel and G. Prasad. *Values of isotropic quadratic forms at S-integral points.* Compositio Math. 83 (1992), no. 3, 347372.

[3] B. Farb and K. Dennis, Noncommutative Algebra, GTM 144, Springer-Verlag, 1993.

[4] G. A. Margulis, G. M. Tomanov, Measure rigidity for almost linear groups and its applications, J. Anal. Math. 69 (1996), 25-54.

[5] S. Mozes, Epimorphic subgroups and invariant measures, Ergodic Theory Dynam. Systems 15 (1995), no. 6, 1207-1210.

[6] D. Poguntke, Normalizers and centralizers of reductive subgroups of almost connected Lie groups, J. Lie Theory 8 (1998), no. 2, 211-217.

[7] M. Ratner, On Raghunathan's measure conjecture. Ann. of Math. (2) 134 (1991), no. 3, 545-607.

[8] R. Shi and B. Weiss, Invariant measures for solvable groups and Diophantine approximation, to appear in Israel J. Math..

[9] J. Tits, Représentations linéaires irréductibles d'un groupe réductif sur un corps quelconque. (French) J. Reine Angew. Math. 247 (1971), 196-220.

[10] G. M. Tomanov, Orbits on homogeneous spaces of arithmetic origin and approximations, in Analysis on homogeneous spaces and representation theory of Lie groups, Okayama-Kyoto (1997), 265-297, Adv. Stud. Pure Math. 26, Math. Soc. Japan, Tokyo, 2000.

[11] C. A. Weibel, An introduction to homological algebra, Cambridge Studies in Advanced Mathematics 38, Cambridge University Press, Cambridge, 1994.

[12] A. Weil, Adeles and algebraic groups, Progress in Mathematics 23, Birkhauser, 1982.

[13] B. Weiss, Divergent trajectories on noncompact parameter spaces, Geom. Funct. Anal. 14 (2004), no. 1, 94-149.

Departement Mathematik, ETH Zürich, Rämistrasse 101, 8092 Zurich, Switzerland

E-mail address: manfred.einsiedler@math.ethz.ch

School of Mathematical Sciences, Xiamen University, Xiamen 361005, PR China

E-mail address: ronggang@xmu.edu.cn